

On the local uniqueness of steady states for the Vlasov-Poisson system

Mikaela Iacobelli *

January 3, 2017

Abstract

Motivated by recent results of Lemou, Méhats, and Raphaël [15] and Lemou [14] concerning the quantitative stability of some suitable steady states for the Vlasov-Poisson system, we investigate the local uniqueness of steady states near these one. This research is inspired by analogous results of Couffrut and Šverák in the context of the 2D Euler equations [6].

1 Introduction

The gravitational Vlasov-Poisson equation modelizes the evolution of a large number of particles subject to their own gravity, under the assumption that both the relativistic effects and the collisions between particles can be neglected. We consider the Vlasov-Poisson system in three dimension:

$$\begin{cases} \partial_t f + v \cdot \nabla_x f - \nabla \phi_f \cdot \nabla_v f = 0, & (t, x, v) \in \mathbb{R}^+ \times \mathbb{R}^3 \times \mathbb{R}^3 \\ f(0, x, v) = f_0(x, v) \geq 0, \quad \int f_0 dx dv = 1. \end{cases} \quad (1.1)$$

where the Newtonian potential ϕ_f is given in terms of the density ρ_f :

$$\rho_f(x) = \int_{\mathbb{R}^3} f(x, v) dv, \quad \text{and} \quad \phi_f(x) = -\frac{1}{4\pi|x|} * \rho_f = K * \rho_f$$

At the beginning of the last century the astrophysicist Sir J. Jeans used this system to model stellar clusters and galaxies [13] and to study their stability properties. In this context it appears in many textbooks on astrophysics such as [4, 10]. In the repulsive case, this system was introduced by A. A. Vlasov around 1937 [21, 22]. Because of the considerable importance in plasma physics and in astrophysics, there is a huge literature on the Vlasov-Poisson system.

The global existence and uniqueness of classical solutions of the Cauchy problem for the Vlasov-Poisson system was obtained by Iordanskii [12] in dimension 1, Ukai-Okabe [20] in the 2-dimensional case, and independently by Lions-Perthame [16] and Pfaffelmoser [18] in the 3-dimensional case (see also [19]). To our knowledge, there are currently no results about existence and uniqueness of classical solutions in dimension greater than 3.

It is important to mention that, parallel to the existence of classical solutions, there have been a considerable amount of work on the existence of weak solutions, in particular

*University of Cambridge, DPMMS Centre for Mathematical Sciences, Wilberforce road, Cambridge CB3 0WB, UK. Email: iacobelli@maths.cam.ac.uk

under very low assumptions on the initial data. We mention in particular the classical result by Arsen'ev [3], who proved global existence of weak solutions under the hypothesis that f_0 is bounded and has finite kinetic energy, and the result of Horst and Hunze [11], where the authors relax the integrability assumption on f_0 . If one wishes to relax even more the integrability assumptions on the initial data then one enters into the framework of the so called renormalized solutions introduced by Di Perna and Lions [7, 8, 9]. The interested reader is referred to the recent papers [1, 5] for more details and references.

One of the main features of the nonlinear transport flow (1.1) is the conservation of the total energy

$$\mathcal{H}(f(t)) = \frac{1}{2} \int_{\mathbb{R}^6} |v|^2 f(t, x, v) \, dx \, dv - \frac{1}{2} \int_{\mathbb{R}^3} |\nabla \phi_f(t, x)|^2 \, dx = \mathcal{H}(f(0)) \quad (1.2)$$

as well as the Casimir functions: for all $G \in C^1([0, \infty], \mathbb{R}^+)$ such that $G(0) = 0$,

$$\int_{\mathbb{R}^6} G(f(t, x, v)) \, dx \, dv = \int_{\mathbb{R}^6} G(f_0(x, v)) \, dx \, dv.$$

1.1 Main result

The goal of this work is to prove a local uniqueness result for steady states of (1.1). In the recent paper [14] (see also [15, 17]), the author proves quantitative stability inequalities for the gravitational Vlasov-Poisson system that will be crucial in the following. More precisely, the author considers a class of steady states $\bar{f}$ to the Vlasov Poisson system, which are decreasing functions of their microscopic energy, and obtains an explicit control of the L^1 distance between $\bar{f}$ and any function f in terms of the energy $\mathcal{H}(f) - \mathcal{H}(\bar{f})$ and the L^1 distance between the rearrangements $\bar{f}^*$ and f^* of $\bar{f}$ and f , respectively.

In the following we give some definition and we state the local functional inequality in [14, Theorem 2]. We first recall the notion of equimeasurability and rearrangement.

Definition 1.1. *Given two integrable nonnegative functions $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$, we say that f and g are equimeasurable if*

$$|\{f > s\}| = |\{g > s\}| \quad \text{for a.e. } s > 0.$$

Then, the (radially decreasing) rearrangement f^* of f is defined as the unique radially decreasing function that is equimeasurable to f . In other words, the level sets of f^* are given by

$$\{f^* > s\} = B_{r(s)}, \quad \text{with } r(s) > 0 \text{ s.t. } |B_{r(s)}| = |\{f > s\}| \text{ for a.e. } s > 0.$$

The following important result is proved in [14, Theorem 2(ii)].

Theorem 1.2. *Consider $\bar{f}$ a compactly supported steady state solution of (1.1) of the form*

$$\bar{f}(x, v) = F(e(x, v)), \quad \text{with } e(x, v) = \frac{|v|^2}{2} + \phi_{\bar{f}}(x), \quad (1.3)$$

where F is a continuous function from $\mathbb{R}$ to $\mathbb{R}^+$ that satisfies the following monotonicity property: there exists $e_0 < 0$ such that $F(e) = 0$ for $e \geq e_0$ and F is a C^1 function on $(-\infty, e_0)$ with $F' < 0$ on $(-\infty, e_0)$. Assume that $f \in L^1 \cap L^\infty(\mathbb{R}^6)$ has finite kinetic energy and is sufficiently close to a translation of $\bar{f}$ in the following sense:

$$\inf_{z \in \mathbb{R}^3} \|\phi_f - \phi_{\bar{f}}(\cdot - z)\|_{L^\infty} + \|\nabla \phi_f - \nabla \phi_{\bar{f}}(\cdot - z)\|_{L^2} < R_0, \quad (1.4)$$

for some suitable constant $R_0 > 0$. Then there exists a constant $K_0 > 0$, depending only on $\bar{f}$, such that

$$\inf_{x_0 \in \mathbb{R}^3} \|f - \bar{f}(\cdot - (x_0, 0))\|_{L^1} \leq \|f^* - \bar{f}^*\|_{L^1} + K_0 [\mathcal{H}(f) - \mathcal{H}(\bar{f}) + \|f^* - \bar{f}^*\|_{L^1}]^{1/2}.$$

where we denote $\bar{f}(\cdot - (x_0, 0)) = \bar{f}(x - x_0, v)$.

An immediate consequence is the following estimate, that will be the starting point of our investigation.

Corollary 1.3. *Let $f, \bar{f}$ be as in Theorem 1.2. Assume in addition that f is equimeasurable to $\bar{f}$. Then*

$$\inf_{x_0 \in \mathbb{R}^3} \|f - \bar{f}(\cdot - (x_0, 0))\|_{L^1}^2 \leq K_0^2 [\mathcal{H}(f) - \mathcal{H}(\bar{f})], \quad (1.5)$$

Let $\bar{f}$ be the stationary solution as above. Our goal is to understand if, nearby $\bar{f}$, there exist other stationary solutions of (1.1). Because stationary solutions of (1.1) correspond to critical points of $\mathcal{H}$ with respect to variations of $\bar{f}$ generated by Hamiltonian flows (see Lemma 2.3 below), it makes sense to consider a “neighborhood” of $\bar{f}$ generated by flows of smooth Hamiltonians. Noticing that $\bar{f}$ is supported in a ball $B_\rho \subset \mathbb{R}^3 \times \mathbb{R}^3$ for some $\rho > 0$ and we shall use the flow of the functions H to move $\bar{f}$, it makes sense to consider Hamiltonians H that are all supported inside $B_{2\rho}$. Hence, one should think of these functions H as the “tangent space” at $\bar{f}$ that will generate the admissible variations.

Let us introduce the following notation:

$$\bar{f}_s^H := (\Phi_s^H)_\# \bar{f} = \bar{f} \circ \Phi_{-s}^H \quad \forall s \in \mathbb{R},$$

where $s \mapsto \Phi_s^H$ is the Hamiltonian flow of H , namely

$$\begin{cases} \partial_s \Phi_s^H = J \nabla H(\Phi_s^H), & J \in \mathbb{R}^{6 \times 6}, \quad J = \begin{pmatrix} 0 & Id \\ -Id & 0 \end{pmatrix} \\ \Phi_0 = Id. \end{cases} \quad (1.6)$$

In other words, $s \mapsto \bar{f}_s^H$ is the variation generated by H , and as H vary this generates a “symplectic” neighborhood of $\bar{f}$. Note that, since $\bar{f}_s^H = \bar{f}_1^{sH}$, to parameterize a neighborhood of $\bar{f}$ it is enough to consider the image of the map¹

$$H \mapsto \bar{f}_1^H. \quad (1.7)$$

We now give some definitions in order to clarify the hypothesis that are needed on the Hamiltonian H . Let us start with the definition of the set $\text{Inv}_{\bar{f}}$ that represents the set of all the Hamiltonians who acts trivially on $\bar{f}$.

Definition 1.4. $\text{Inv}_{\bar{f}} := \{H \in C^{1,1}(\mathbb{R}^6) : \{H, \bar{f}\} = 0\}$

This definition is motivated by the following simple result:

¹This resembles to the exponential map in Riemannian geometry, where a neighborhood of a point $x \in M$ is obtained as the image of a neighborhood of 0 in $T_x M$ via the map

$$v \mapsto \gamma_v(1),$$

where $s \mapsto \gamma_v(s)$ is the geodesic starting at x with velocity v .

Lemma 1.5. *If $H \in \text{Inv}_{\bar{f}}$ then $(\Phi_t^H)_\# \bar{f} = \bar{f}$, i.e., the Hamiltonian flow of H does not move $\bar{f}$.*

Proof. The function $\bar{f}_s^H := (\Phi_s^H)_\# \bar{f}$ solves the transport equation

$$\partial_s \bar{f}_s^H + \text{div}(J\nabla H \bar{f}_s^H) = 0, \quad \bar{f}_s^H|_{s=0} = \bar{f}.$$

Since also $\bar{f}$ is a solution to this equation (because $\text{div}(J\nabla H \bar{f}) = \{H, \bar{f}\} = 0$) and the vector field $J\nabla H$ is Lipschitz, $\bar{f}_s \equiv \bar{f}$ by uniqueness for the above transport equation. $\square$

It follows by the lemma above that if H belongs to $\text{Inv}_{\bar{f}}$ then Φ_s^H is not moving $\bar{f}$. Since our goal is to use Hamiltonians H to parameterize a neighborhood of $\bar{f}$, there is no reason to consider H that belong to $\text{Inv}(\bar{f})$, and it make sense to exclude them. Actually, for some technical reasons that will be more clear later, we shall need to impose a quantitative version of the condition $H \notin \text{Inv}_{\bar{f}}$. To do that, we introduce the family of sets

$$A_k := \{H \neq 0 : \|\nabla H\|_{L^1} \leq k\|\{H, \bar{f}\}\|_{L^1}\}, \quad k \geq 1.$$

Remark 1.6. *We note that $\text{Inv}_{\bar{f}} = \bigcap_{k \in \mathbf{N}} A_k^c$. Indeed, if $H \in \bigcap_{k \in \mathbf{N}} A_k^c$ then $\|\nabla H\|_{L^1}/k \geq \|\{H, \bar{f}\}\|_{L^1}$ for all $k \in \mathbf{N}$. Thus $\{H, \bar{f}\} \equiv 0$, which implies that $H \in \text{Inv}_{\bar{f}}$. Viceversa, if $H \in \text{Inv}_{\bar{f}}$ then clearly $H \in \bigcap_{k \in \mathbf{N}} A_k^c$.*

Because of this observation, we see that

$$H \notin \text{Inv}_{\bar{f}} \iff \exists k \text{ such that } H \in A_k.$$

Motivated by this fact, in the sequel we shall fix k and consider only Hamiltonians that belong to A_k . Of course this is more restrictive than assuming only $H \notin \text{Inv}_{\bar{f}}$ but at the moment it is not clear to us how to remove such an assumption.

Going further in our preliminary analysis, we observe that all translations of $\bar{f}$ are trivially stationary solutions. However, translations in v are automatically controlled by the kinetic energy and indeed they do not appear in (1.5). To “kill” the space of translations in x , we will assume that $\text{Bar}_x(\bar{f}_1^H) = \text{Bar}_x(\bar{f})$, where

$$\text{Bar}_x(f) := \int_{\mathbb{R}^6} x f(x, v) dx dv \in \mathbb{R}^3$$

denotes the “barycenter (in x)” of f . We want to emphasize that this is not a restrictive assumption on H , since one could remove it by adding to H a Hamiltonian corresponding to translations in the x variable in order to recenter the barycenter of f . Since this would not add major technical difficulties to the proof but may distract the reader from the essential points, we decided to impose this barycenter condition on $\bar{f}_1^H$.

As a final consideration, since our goal is prove that there are no steady states to (1.1) in a neighborhood of $\bar{f}$ generated via the map (1.7), we shall need to assume that our Hamiltonians H are small in some suitable topology.

Our main theorem asserts that, for Hamiltonians small enough in a sufficiently strong Sobolev norm that are quantitatively away from $\text{Inv}_{\bar{f}}$, there cannot be a stationary point of the form $\bar{f}_1^H$.

Theorem 1.7. *Let $\bar{f}$ be as in (1.3), where F is a continuous function from $\mathbb{R}$ to $\mathbb{R}^+$ that satisfies the following monotonicity property: there exists $e_0 < 0$ such that $F(e) = 0$ for $e \geq e_0$ and F is a C^1 function on $(-\infty, e_0)$ with $F' < 0$ on $(-\infty, e_0)$. Let $\rho > 0$ be such that $\text{supp}(\bar{f}) \subset B_\rho$. Also, assume that $\bar{f} \in W^{2,q}(\mathbb{R}^6)$ for some $q > 3$. Then the following local uniqueness result for steady states holds:*

Let $r \geq 22$, and given $\rho, \varepsilon, k > 0$ consider the space of functions

$$N_\varepsilon^k := \{\bar{f}_1^H : \text{supp}(H) \subset B_{2\rho}, \text{Bar}_x(\bar{f}_1^H) = \text{Bar}_x(\bar{f}), H \in A_k, \|H\|_{W^{r,2}} \leq \varepsilon\}.$$

Then, fixed $k \in \mathbb{N}$, there is no stationary state for (1.1) in N_ε^k for ε small enough.

1.1.1 Comments

Starting from the seminal paper of Arnold about the geometric interpretation of the Euler equations as L^2 -geodesics in the space of measure preserving diffeomorphisms [2], Choffrut and Šverák recently obtained a related result for the 2D Euler equation [6]. The basic idea there is that, under the evolution given by the 2D incompressible Euler equations, the vorticity is transported by an incompressible vector field, hence the measure of all its super-level sets is constant. This means that, given an initial vorticity ω_0 , its evolution $\omega(t)$ is in the same equimeasurability class of ω_0 . This allows one to foliate the space of vorticities into a family of leaves $\mathcal{O}_{\omega_0}$ (the equimeasurability class of ω_0), and the Euler equations preserve these leaves. In addition, thanks to the Hamiltonian structure of the Euler equations, one can characterize stationary solutions as critical points of the Hamiltonian energy $\mathcal{E}$ restricted to the orbits.

In other words, one has the following situation: the space of vorticities is foliated by the orbits $\mathcal{O}_\omega$, and the equilibria are the critical points of $\mathcal{E}$ restricted to the orbits. In finite dimension, the implicit function theorem would give the following: if $\mathcal{O}_\omega$ is smooth near a point $\bar{\omega} \in \mathcal{O}_\omega$, and if $\bar{\omega}$ is a non-degenerate critical point of $\mathcal{E}$ in $\mathcal{O}_\omega$, then near $\bar{\omega}$ the set of equilibria form a smooth manifold transversal to the foliation. In addition, the dimension of this manifold is equation to the co-dimension of the orbits. In particular, in a non-degenerate situation, the equilibria are locally in one-to-one correspondence with the orbits. In [6] the authors obtain an analogue of this correspondence in the infinite dimensional context of Euler equations. There, the authors use an infinite dimensional version of the implicit function theorem in the space of C^∞ function, via a Nash-Moser's iteration.

With respect to their result, here we have different assumptions and results. These are motivated by the following:

- Since the Vlasov-Poisson system (1.1) is Hamiltonian, given an initial condition f_0 its evolution f_t under the Vlasov-Poisson system will also be in the same equimeasurability class. However, while Hamiltonian maps and measure preserving maps coincide in 2-dimension, they are very different in higher dimension (for instance, Hamiltonian maps preserve the symplectic structure). Because solutions to the 3D Vlasov-Poisson systems describe a Hamiltonian evolution of particles in the phase-space $\mathbb{R}^3 \times \mathbb{R}^3$, there is no natural reason in this context why there should be only one stationary state in the same equimeasurability class. In particular, as already observed before, stationary solutions of (1.1) correspond to critical points of $\mathcal{H}$ with respect to variations of $\bar{f}$ generated by Hamiltonian flows, and not with respect to arbitrary measure preserving variations. This is why we need to look at functions f that can be connected to $\bar{f}$ via a Hamiltonian flow, namely $f = \bar{f}_1^H$ for some H .

- The smallness assumption on $\|\nabla^r H\|_{L^2}$ is natural, and actually weaker than the one in [6], since smallness there is measured in the C^∞ topology.
- As already mentioned before, the assumption on $\text{Bar}_x(\bar{f}_1^H)$ is not fundamental: one could easily remove it by replacing it with $H - H_0$, where H_0 corresponds to a translation in the phase space (multiplied by a suitable cut-off function, to make it compactly supported). What is more essential is our assumption $H \in A_k$, and it is unclear at the moment how to remove it. It is our plan to address this issue in a future work.

The goal of the next section is to prove our main theorem.

2 Proof of the Theorem 1.7

2.1 Strategy of the proof

The idea of the proof is the following: first, by exploiting the results in [14], we prove that if $\bar{f}_1^H$ has the same barycenter as $\bar{f}$, then

$$\|\bar{f}_1^H - \bar{f}\|_{L^1}^2 \leq K_0^2[\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})].$$

Secondly we show that if $\bar{f}_1^H$ is stationary, by a Taylor expansion of $s \mapsto \mathcal{H}(\bar{f}_s^H)$ we can prove that

$$\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f}) \leq C\|\nabla H\|_X^3$$

for some suitable norm $\|\cdot\|_X$ of ∇H .

Combining these two estimates, we get

$$\|\bar{f}_1^H - \bar{f}\|_{L^1} \leq C\|\nabla H\|_X^{3/2}.$$

We then relate the two quantities appearing in the above expression: more precisely, we first show that

$$\|\bar{f}_1^H - \bar{f}\|_{L^1} \approx \|\{H, \bar{f}\}\|_{L^1} = \|\nabla H \cdot J\nabla \bar{f}\|_{L^1},$$

and then we use our quantitative assumption on the fact that H does not belong to $\text{Inv}_{\bar{f}}$ (namely, $H \in A_k$) to say that

$$\|\{H, \bar{f}\}\|_{L^1} \approx \|\nabla H\|_{L^1}.$$

In this way we get

$$\|\nabla H\|_{L^1} \leq C\|\nabla H\|_X^{3/2}.$$

Finally, exploiting the smallness $\|\nabla^r H\|_{L^2} \leq \varepsilon$ and interpolation estimates, we are able to relate the two norms above and conclude that

$$\|\nabla H\|_{L^2} \leq C\|\nabla H\|_{L^2}^{1+\delta}$$

for some $\delta > 0$, which yields a contradiction when $\|\nabla H\|_{L^2}$ is small enough.

2.2 Lower bound

We recall the definition of the baricenter of f :

$$\text{Bar}_x(f) = \int_{\mathbb{R}^6} x f(x, v) dx dv.$$

Lemma 2.1. *Let $\bar{f}$ be as in (1.3), where F is a continuous function from $\mathbb{R}$ to $\mathbb{R}^+$ that satisfies the following monotonicity property: there exists $e_0 < 0$ such that $F(e) = 0$ for $e \geq e_0$ and F is a C^1 function on $(-\infty, e_0)$ with $F' < 0$ on $(-\infty, e_0)$. Let $H \in C^2(\mathbb{R}^6)$ and consider the function $\bar{f}_1^H := (\Phi_1^H)_\# \bar{f}$ for some $H \in C^{1,1}$ with $\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} \leq \eta$. Also, assume that*

$$\text{Bar}_x(\bar{f}) = \text{Bar}_x(\bar{f}_1^H). \quad (2.1)$$

Then, if η is small enough,

$$\|\bar{f}_1^H - \bar{f}\|_{L^1}^2 \leq \hat{K}_0[\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})], \quad (2.2)$$

where $\hat{K}_0$ depends on the diameter of the support of $\bar{f}$ and $\bar{f}_1^H$.

Proof. Note that, if η is small enough, the function $f = \bar{f}_1^H$ satisfies (1.4). Let $(x_0, 0)$ be the point where the minimum is achieved in (1.5). By definition,

$$\begin{aligned} \text{Bar}_x(\bar{f}_1^H(\cdot - (x_0, 0))) &= \int_{\mathbb{R}^6} x \bar{f}_1^H(x - x_0, v) dx dv \\ &= \int_{\mathbb{R}^6} (x - x_0) \bar{f}_1^H(x - x_0, v) dx dv + x_0, \\ &= \text{Bar}_x(\bar{f}_1^H) + x_0. \end{aligned}$$

hence

$$\begin{aligned} |x_0| &= |\text{Bar}_x(\bar{f}_1^H) - \text{Bar}_x(\bar{f}_1^H(\cdot - (x_0, 0)))| \stackrel{(2.1)}{=} |\text{Bar}_x(\bar{f}) - \text{Bar}_x(\bar{f}_1^H(\cdot - (x_0, 0)))| \\ &\leq \int_{\mathbb{R}^6} |x| |\bar{f} - \bar{f}_1^H(\cdot - (x_0, v_0))| dx dz \\ &\leq C \|\bar{f} - \bar{f}_1^H(\cdot - (x_0, v_0))\|_{L^1} \leq C[\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})]^{1/2}, \end{aligned}$$

where we used that $\bar{f}$ and $\bar{f}_1^H$ are compactly supported, so $|(x, v)|$ is bounded on the support of $\bar{f}$ and $\bar{f}_1^H(\cdot - (x_0, v_0))$. Thus,

$$\begin{aligned} \|\bar{f} - \bar{f}_1^H\|_{L^1} &\leq \|\bar{f} - \bar{f}(\cdot + (x_0, v_0))\|_{L^1} + \|\bar{f}(\cdot + (x_0, v_0)) - \bar{f}_1^H\|_{L^1} \\ &\leq |x_0| \|\nabla \bar{f}\|_{L^1} + [\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})]^{1/2} \leq C[\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})]^{1/2}, \end{aligned}$$

which concludes the proof. $\square$

2.3 Upper bound

The aim of this section is to provide an estimate of the difference between the energy of $\bar{f}$ and of $\bar{f}_1^H$ in terms H , under the additional assumption that $\bar{f}_1^H$ is a stationary solution for (1.1). More precisely, we prove the following:

Proposition 2.2. *Let $\bar{f}$ be a compactly supported steady state such that $\bar{f} \in L^\infty(\mathbb{R}^6)$, and that $\bar{f} \in W^{2,q}(\mathbb{R}^6)$ for some $q > 3$. Let $H \in C^2(\mathbb{R}^6)$. Also, assume that $\bar{f}_1^H = (\Phi_1^H)_\# \bar{f}$ is a stationary solution for (1.1). Then the following estimate holds:*

$$|\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})| \leq C \|\nabla H\|_{L^\infty} \left(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} \right)^2,$$

where C is a constant depending only on $\bar{f}$.

As a first step towards the proof of the above result, we aim to give a characterization of the stationary solutions of (1.1) in terms of the energy of the system $\mathcal{H}$.

Lemma 2.3. *Let $f : \mathbb{R}^6 \rightarrow \mathbb{R}$ be a compactly supported function. Then f is a steady state for (1.1) if and only if*

$$\frac{d}{ds} \mathcal{H}(f_s^H)|_{s=0} = 0$$

for all $H \in C^2(\mathbb{R}^6)$, where $f_s^H := (\Phi_s^H)_\# f$.

Proof. Fix $H \in C^2(\mathbb{R}^6)$, and consider its flow Φ_s^H . To simplify the notation we set $\Phi_s^H = \Phi_s$. Also, it will be convenient to write $\Phi_s = (\Phi_s^x, \Phi_s^v) : \mathbb{R}^6 \rightarrow \mathbb{R}^3 \times \mathbb{R}^3$.

Given a compactly supported function f , we compute the first variation of the Hamiltonian $\mathcal{H}$ around f along f_s^H :

$$\begin{aligned} \frac{d}{ds} \mathcal{H}(f_s^H) &= \frac{d}{ds} \left[\frac{1}{2} \int_{\mathbb{R}^6} |v|^2 f_s^H(x, v) \, dx \, dv - \frac{1}{2} \int_{\mathbb{R}^6 \times \mathbb{R}^6} K(x - y) f_s^H(x, v) f_s^H(y, w) \, dx \, dv \, dy \, dw \right] \\ &= \frac{d}{ds} \left[\frac{1}{2} \int_{\mathbb{R}^6} |\Phi_s^v(x, v)|^2 f(x, v) \, dx \, dv \right. \\ &\quad \left. - \frac{1}{2} \int_{\mathbb{R}^6 \times \mathbb{R}^6} K(\Phi_s^x(x, v) - \Phi_s^x(y, v)) f(x, v) f(y, w) \, dx \, dv \, dy \, dw \right] \\ &= \int_{\mathbb{R}^6} \Phi_s^v(x, v) \partial_s \Phi_s^v(x, v) f(x, v) \, dx \, dv \\ &\quad - \frac{1}{2} \left[\int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(\Phi_s^x(x, v) - \Phi_s^x(y, v)) \cdot \right. \\ &\quad \left. \cdot \partial_s (\Phi_s^x(x, v) - \Phi_s^x(y, v)) f(x, v) f(y, w) \, dx \, dv \, dy \, dw \right]. \end{aligned}$$

Recalling that

$$\partial_s \Phi_s = (\nabla_v H(\Phi_s), -\nabla_x H(\Phi_s)),$$

we have

$$\begin{aligned} \frac{d}{ds} \mathcal{H}(f_s^H) &= - \int_{\mathbb{R}^6} \Phi_s^v(x, v) \cdot \nabla_x H(\Phi_s(x, v)) f(x, v) \, dx \, dv \\ &\quad - \frac{1}{2} \left[\int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(\Phi_s^x(x, v) - \Phi_s^x(y, v)) \cdot \right. \\ &\quad \left. \cdot (\nabla_v H(\Phi_s(x, v)) - \nabla_v H(\Phi_s(y, v))) f(x, v) f(y, w) \, dx \, dv \, dy \, dw \right]. \end{aligned}$$

Also, since $K(x - y) = K(y - x)$, we see that $\nabla_x K(x - y) = -\nabla_x K(y - x)$, so we can rewrite the above expression as

$$\begin{aligned} \frac{d}{ds} \mathcal{H}(f_s^H) &= - \int_{\mathbb{R}^6} \Phi_s^v(x, v) \cdot \nabla_x H(\Phi_s(x, v)) f(x, v) \, dx \, dv \\ &\quad - \int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(\Phi_s^x(x, v) - \Phi_s^x(y, v)) \cdot \nabla_v H(\Phi_s(x, v)) f(x, v) f(y, w) \, dx \, dv \, dy \, dw. \end{aligned}$$

Also, using that Φ_s preserves the Lebesgue measure and that $\Phi_s^{-1} = \Phi_{-s}$, we can rewrite the first variation in the following way:

$$\begin{aligned} \frac{d}{ds} \mathcal{H}(f_s^H) &= - \int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) f(\Phi_{-s}(x, v)) \, dx \, dv \\ &\quad - \int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(x - y) \cdot \nabla_v H(x, v) f(\Phi_{-s}(x, v)) f(\Phi_{-s}(y, w)) \, dx \, dv \, dy \, dw. \end{aligned} \quad (2.3)$$

In particular, since $\Phi_s = Id$ for $s = 0$, we see that

$$\begin{aligned} \frac{d}{ds} \mathcal{H}(f_s)|_{s=0} &= - \int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) f(x, v) \, dx \, dv \\ &\quad - \int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(x - y) \cdot \nabla_v H(x, y) f(x, v) f(y, w) \, dx \, dv \, dy \, dw. \end{aligned} \quad (2.4)$$

On the other hand, for f to be a stationary solution for the system (1.1) means that

$$\operatorname{div}_x(vf(x, v)) - \operatorname{div}_v(\nabla \phi_f(x) f(x, v)) = 0,$$

or equivalently, that for all $\psi \in C^1$,

$$- \int_{\mathbb{R}^6} v \cdot \nabla_x \psi(x, v) f(x, v) \, dx \, dv + \int_{\mathbb{R}^3} \nabla_x \phi_f(x) \cdot \nabla_v \psi(x, v) f(x, v) \, dx \, dv = 0. \quad (2.5)$$

Since

$$\int \nabla_x K(x - y) f(y, w) \, dy \, dw = -\nabla \phi_f,$$

(2.4) proves that

$$\frac{d}{ds} \mathcal{H}(f_s^H)|_{s=0} = 0 \quad \Longleftrightarrow \quad (2.5) \text{ holds with } \psi = H.$$

Since C^2 functions are dense in C^1 for the C^1 topology, this proves the result. $\square$

2.3.1 Second variation for $\mathcal{H}$

As a second step, we compute the second variation for $\mathcal{H}$, in line with the computation of the first variation (2.3). Here we consider as initial condition $\bar{f}$ and, given a Hamiltonian $H \in C^2$, we consider $\bar{f}_s^H := \bar{f} \circ \Phi_{-s}^H$. As before, to simplify the notation, we set $\Phi_s = \Phi_s^H$. Also, we define

$$g := \nabla \bar{f} \cdot J \nabla H = \{H, \bar{f}\} \quad (2.6)$$

and we observe that

$$\frac{d}{ds} \bar{f}(\Phi_{-s}) = g(\Phi_{-s}). \quad (2.7)$$

Using the equations (2.3) and (2.7) the second variation is given by the following:

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) &= \left[\int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) g(\Phi_{-s}(x, v)) \, dx \, dv \right] \\ &\quad + \left[\int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(x - y) \cdot \nabla_v H(x, v) g(\Phi_{-s}(x, v)) \bar{f}(\Phi_{-s}(y, w)) \, dx \, dv \, dy \, dw \right] \\ &\quad + \left[\int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(x - y) \cdot \nabla_v H(x, v) \bar{f}(\Phi_{-s}(x, v)) g(\Phi_{-s}(y, w)) \, dx \, dv \, dy \, dw \right]. \end{aligned}$$

Thus, we obtain:

$$\begin{aligned} \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) &= \left[\int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) g(\Phi_{-s}(x, v)) dx dv \right] \\ &+ \left[\int_{\mathbb{R}^6 \times \mathbb{R}^6} \nabla_x K(x - y) \cdot [\nabla_v H(x, v) - \nabla_w H(y, w)] \bar{f}(\Phi_{-s}(x, v)) g(\Phi_{-s}(y, w)) dx dv dy dw \right]. \end{aligned} \quad (2.8)$$

2.3.2 Proof of Proposition 2.2

As in the previous section, we set $\bar{f}_s^H := \bar{f} \circ \Phi_{-s}^H$ and $\Phi_s := \Phi_s^H$. Recall that, by assumption $\bar{f}_1^H$, is a stationary solution of (1.1).

We now study the Taylor expansion of the Hamiltonian of the gravitational Vlasov Poisson system both in $\bar{f}$ and in $\bar{f}_1^H$. Since $\bar{f}$ and $\bar{f}_1^H$ are two stationary solutions, it follows by Lemma 2.3 applied both to $\bar{f}$ and to $\bar{f}_1^H$ that

$$\frac{d}{ds} \mathcal{H}(\bar{f}_s^H)|_{s=0} = \frac{d}{ds} \mathcal{H}(\bar{f} \circ \Phi_{-s})|_{s=0} = 0$$

and

$$\frac{d}{ds} \mathcal{H}(\bar{f}_s^H)|_{s=1} = \frac{d}{d\tau} \mathcal{H}(\bar{f}_1^H \circ \Phi_{-\tau})|_{\tau=0} = 0.$$

Hence, by Taylor's formula,

$$\mathcal{H}(\bar{f}_1^H) = \mathcal{H}(\bar{f}) + \int_0^1 (1-s) \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) ds$$

and

$$\mathcal{H}(\bar{f}) = \mathcal{H}(\bar{f}_1^H) + \int_0^1 s \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) ds.$$

Therefore,

$$\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f}) = \int_0^1 (1-2s) \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) ds.$$

Since

$$\int_0^1 (1-2s) ds = 0$$

we can add a constant term in the integral, and we get

$$\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f}) = \frac{1}{2} \int_0^1 (1-2s) \left(\frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) - \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H)|_{s=0} \right) ds. \quad (2.9)$$

Thanks to the latter computation, in order to estimate the left hand side of (2.9), we can estimate

$$\frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H) - \frac{d^2}{ds^2} \mathcal{H}(\bar{f}_s^H)|_{s=0}$$

in terms of the regularity of the Hamiltonian H , and of $\bar{f}$. Recalling that $\Phi_0 = \text{Id}$, we have the following expression:

$$\begin{aligned}
& \frac{d^2}{d^2 s} \mathcal{H}(\bar{f}_s^H) - \frac{d^2}{d^2 s} \mathcal{H}(\bar{f}_s^H)|_{s=0} = \int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) g(\Phi_{-s}(x, v)) \, dx \, dv \\
& - \int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) g(x, v) \, dx \, dv \\
& + \int_{\mathbb{R}^6 \times 6} \nabla_x K(x - y) \cdot [\nabla_v H(x, v) - \nabla_w H(y, w)] \bar{f}(\Phi_{-s}(x, v)) g(\Phi_{-s}(y, w)) \, dx \, dv \, dy \, dw \\
& - \int_{\mathbb{R}^6 \times 6} \nabla_x K(x - y) \cdot [\nabla_v H(x, v) - \nabla_w H(y, w)] \bar{f}(x, v) g(y, w) \, dx \, dv \, dy \, dw. \\
& \leq T_1 + T_2,
\end{aligned} \tag{2.10}$$

where

$$\begin{aligned}
T_1 &:= \left| \int_{\mathbb{R}^6} v \cdot \nabla_x H(x, v) [g(\Phi_{-s}(x, v)) - g(x, v)] \, dx \, dv \right|, \\
T_2 &:= \left| \int_{\mathbb{R}^6 \times 6} \nabla_x K(x - y) \cdot [\nabla_v H(x, v) - \nabla_w H(y, w)] \bar{f}(\Phi_{-s}(x, v)) g(\Phi_{-s}(y, w)) \, dx \, dv \, dy \, dw \right. \\
&\quad \left. - \int_{\mathbb{R}^6 \times 6} \nabla_x K(x - y) \cdot [\nabla_v H(x, v) - \nabla_w H(y, w)] \bar{f}(x, v) g(y, w) \, dx \, dv \, dy \, dw \right|.
\end{aligned}$$

We begin by controlling T_1 .

By the Fundamental Theorem of Calculus we have that

$$\begin{aligned}
T_1 &\leq C \|\nabla_x H\|_{L^\infty} \int_{\mathbb{R}^6} \left| g(\Phi_{-s}(x, v)) - g(x, v) \right| \, dx \, dv \\
&\leq C \|\nabla_x H\|_{L^\infty} \int_{\mathbb{R}^6} \left(\int_0^1 \left| \nabla g(\Phi_{-\tau s}(x, v)) \cdot \partial_\tau \Phi_{-\tau s}(x, v) \right| \, d\tau \right) \, dx \, dv
\end{aligned}$$

Using that $\partial_s \Phi_s = J \nabla H(\Phi_s)$ and that Φ_s preserves the volumes, we get

$$\begin{aligned}
T_1 &\leq C \|\nabla H\|_\infty^2 \int_{\mathbb{R}^6} \left(\int_0^1 \left| \nabla g(\Phi_{-\tau s}(x, v)) \right| \, d\tau \right) \, dx \, dv \\
&= C \|\nabla H\|_\infty^2 \int_{\mathbb{R}^6} \left(\int_0^1 \left| \nabla g(x, v) \right| \, ds \right) \, dx \, dv \\
&\leq C \|\nabla H\|_\infty^2 \|\nabla g\|_{L^1}.
\end{aligned}$$

Thus,

$$T_1 \leq C \|\nabla H\|_\infty^2 \|\nabla g\|_{L^1}.$$

By the definition of g in (2.6), we have that

$$\nabla g = \nabla^2 H \cdot J \nabla \bar{f} + \nabla H \cdot J \nabla^2 \bar{f}. \tag{2.11}$$

Therefore,

$$\|\nabla g\|_{L^1} \leq \|\nabla^2 H\|_{L^\infty} \|\nabla \bar{f}\|_{L^1} + \|\nabla H\|_{L^\infty} \|\nabla^2 \bar{f}\|_{L^1} \leq C (\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty}),$$

where C depends on $\|\nabla \bar{f}\|_{L^1}$ and $\|\nabla^2 \bar{f}\|_{L^1}$. In conclusion, the first term T_1 can be estimate as follows:

$$T_1 \leq C \|\nabla H\|_{L^\infty}^2 \left(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} \right). \quad (2.12)$$

We now estimate the second term:

$$T_2 \leq \|\nabla H\|_{L^\infty} \int_{\mathbb{R}^{6 \times 6}} |\nabla_x K(x-y)| |\bar{f}(\Phi_{-s}(x,v))g(\Phi_{-s}(y,w)) - \bar{f}(x,v)g(y,w)| dx dv dy dw.$$

Adding and subtracting $\bar{f}(x,v)g(\Phi_{-s}(y,w))$, we can bound

$$\begin{aligned} T_2 &\leq \|\nabla H\|_{L^\infty} \int_{\mathbb{R}^{6 \times 6}} |\nabla_x K(x-y)| |\bar{f}(\Phi_{-s}(x,v))g(\Phi_{-s}(y,w)) - \bar{f}(x,v)g(\Phi_{-s}(y,w))| dx dv dy dw \\ &\quad + \|\nabla H\|_{L^\infty} \int_{\mathbb{R}^{6 \times 6}} |\nabla_x K(x-y)| |\bar{f}(x,v)g(\Phi_{-s}(y,w)) - \bar{f}(x,v)g(y,w)| dx dv dy dw \\ &\leq C \|\nabla H\|_{L^\infty} \|g\|_{L^\infty} \int_{\mathbb{R}^3 \times B_R} |\nabla_x K(x-y)| |\bar{f}(\Phi_{-s}(x,v)) - \bar{f}(x,v)| dx dv dy dw \\ &\quad + C \|\nabla H\|_{L^\infty} \|\bar{f}\|_{L^\infty} \int_{B_R \times \mathbb{R}^3} |\nabla_x K(x-y)| |g(\Phi_{-s}(y,w)) - g(y,w)| dx dv dy dw. \end{aligned}$$

Using as before the Fundamental Theorem of Calculus and the fact that Φ_s is measure preserving, we have

$$\begin{aligned} T_2 &\leq C \|\nabla H\|_{L^\infty} \|g\|_{L^\infty} \int_{\mathbb{R}^6 \times B_R} |\nabla_x K(x-y)| \cdot \left(\int_0^1 |\nabla \bar{f}(\Phi_{-\tau s}(x,v)) \cdot \partial_s \Phi_{-\tau s}(x,v)| d\tau \right) dx dv dy dw \\ &\quad + C \|\nabla H\|_{L^\infty} \|\bar{f}\|_{L^\infty} \int_{B_R \times \mathbb{R}^6} |\nabla_x K(x-y)| \cdot \left(\int_0^1 |\nabla g(\Phi_{-\tau s}(y,w)) \cdot \partial_s \Phi_{-\tau s}(y,w)| d\tau \right) dx dv dy dw. \end{aligned}$$

By the definition of g , (2.6) we obtain

$$\begin{aligned} T_2 &\leq C \|\nabla H\|_{L^\infty} \|g\|_{L^\infty} \int_{\mathbb{R}^6 \times B_R} |\nabla_x K(x-y)| \left(\int_0^1 |g(\Phi_{-\tau s}(x,v))| d\tau \right) dx dv dy dw \\ &\quad + C \|\nabla H\|_{L^\infty} \|\bar{f}\|_{L^\infty} \int_{B_R \times \mathbb{R}^6} |\nabla_x K(x-y)| \cdot \left(\int_0^1 |\nabla g(\Phi_{-\tau s}(y,w)) \cdot J\nabla H(\Phi_{-\tau s})(-s)| d\tau \right) dx dv dy dw, \end{aligned}$$

and using Hölder inequality we get

$$\begin{aligned} T_2 &\leq C \|\nabla H\|_{L^\infty} \|g\|_{L^\infty}^2 \int_{B_R \times B_R} |\nabla_x K(x-y)| dx dy \\ &\quad + C \|\nabla H\|_{L^\infty}^2 \|\bar{f}\|_{L^\infty} \|\nabla g\|_{L^q} \left(\int_{B_R \times B_R} |\nabla_x K(x-y)|^p dx dy \right)^{\frac{1}{p}}, \end{aligned}$$

where p and q are conjugate exponents. In order to have integrability of the gradient of the kernel K , we need $p < \frac{3}{2}$. Therefore, in the previous estimates we need to assume that $\|\nabla g\|_{L^q}$ is finite for some $q > 3$. Thus

$$T_2 \leq C(\|\nabla H\|_{L^\infty} \|g\|_{L^\infty}^2 + \|\nabla H\|_{L^\infty}^2 \|\nabla g\|_{L^q}), \quad q > 3,$$

where C depends on $\|\bar{f}\|_{L^\infty}$. As in the estimate of the term T_1 we use (2.6) and (2.11) to get

$$\|g\|_{L^\infty} \leq \|\nabla H\|_{L^\infty} \|\nabla \bar{f}\|_{L^\infty} \leq C \|\nabla H\|_{L^\infty}$$

and

$$\|\nabla g\|_{L^q} \leq \|\nabla^2 H\|_{L^\infty} \|\nabla \bar{f}\|_{L^q} + \|\nabla H\|_{L^\infty} \|\nabla^2 \bar{f}\|_{L^q} \leq C(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty}),$$

where C depends on $\|\nabla \bar{f}\|_{L^q}$ and $\|\nabla^2 \bar{f}\|_{L^q}$ for some $q > 3$. Since by assumption $\bar{f} \in W^{2,q}(\mathbb{R}^6)$ for some $q > 3$, we have prove that

$$T_2 \leq C(\|\nabla H\|_{L^\infty}^3 + \|\nabla H\|_{L^\infty}^2 \|\nabla^2 H\|_{L^\infty}). \quad (2.13)$$

Hence, combining (2.10), (2.12), and (2.13) we get

$$|\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})| \leq C(\|\nabla H\|_{L^\infty}^3 + \|\nabla H\|_{L^\infty}^2 \|\nabla^2 H\|_{L^\infty}),$$

where C depends only on $\|\bar{f}\|_{L^\infty}$ and on $\|\nabla \bar{f}\|_{L^q}$ and $\|\nabla^2 \bar{f}\|_{L^q}$, for some $q > 3$. $\square$

2.4 Comparing $\|\bar{f} - \bar{f}_1^H\|_{L^1}$ and $\|g\|_{L^1}$

Lemma 2.4. *Let $\bar{f}$ be a compactly supported steady state such that $\nabla \bar{f}, \nabla^2 \bar{f} \in L^1(\mathbb{R}^6)$. Also, let $H \in C^2(\mathbb{R}^6)$, and define g as in (2.6). Set $\bar{f}_1^H := \bar{f} \circ \Phi_{-1}^H$. Then*

$$\left| \|\bar{f} - \bar{f}_1^H\|_{L^1} - \|g\|_{L^1} \right| \leq C \|\nabla H\|_{L^\infty} \left(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} e^{\|\nabla^2 H\|_{L^\infty}} \right),$$

where C depends only on $\bar{f}$.

Proof. Set $\bar{f}_s^H := \bar{f} \circ \Phi_{-s}^H$. Then, by the definition of the flow Φ_s (see (1.6)), we deduce

$$\partial_s \bar{f}_s^H = -J \nabla H \cdot \nabla \bar{f}_s^H, \quad \partial_s \bar{f}_s^H|_{s=0} = -J \nabla H \cdot \nabla \bar{f}, \quad (2.14)$$

therefore

$$\begin{aligned} \bar{f}_1^H - \bar{f} &= \int_0^1 \partial_s \bar{f}_s^H ds = \partial_s \bar{f}_s^H|_{s=0} + \int_0^1 (\partial_s \bar{f}_s^H - \partial_s \bar{f}_s^H|_{s=0}) ds \\ &\stackrel{(2.14)}{=} -J \nabla H \cdot \nabla \bar{f} + \int_0^1 (\partial_s \bar{f}_s^H - \partial_s \bar{f}_s^H|_{s=0}) ds \\ &\stackrel{(2.6)}{=} -g + \int_0^1 (\partial_s \bar{f}_s^H - \partial_s \bar{f}_s^H|_{s=0}) ds. \end{aligned}$$

Thus,

$$\left| \int_{\mathbb{R}^6} |\bar{f}_1^H - \bar{f}| dx dv - \int_{\mathbb{R}^6} |g| dx dv \right| \leq \int_0^1 \int_{\mathbb{R}^6} |\partial_s \bar{f}_s^H - \partial_s \bar{f}_s^H|_{s=0}| dx dv ds.$$

Using the equations (2.14), we have:

$$\begin{aligned} \left| \int_{\mathbb{R}^6} |\bar{f}_1^H - \bar{f}| dx dv - \int_{\mathbb{R}^6} |g| dx dv \right| &\leq \int_0^1 \int_{\mathbb{R}^6} |J\nabla H \cdot \nabla \bar{f}_s^H - J\nabla H \cdot \nabla \bar{f}| dx dv ds \\ &\leq \int_0^1 ds \int_{\mathbb{R}^6} |\nabla H| |\nabla \bar{f} \circ \Phi_{-s}^H \cdot \nabla \Phi_{-s}^H - \nabla \bar{f}| dx dv. \end{aligned}$$

Adding and subtracting $\nabla \bar{f} \circ \Phi_{-s}^H$, this gives

$$\begin{aligned} \left| \int_{\mathbb{R}^6} |\bar{f}_1^H - \bar{f}| dx dv - \int_{\mathbb{R}^6} |g| dx dv \right| &\leq \|\nabla H\|_{L^\infty} \left(\int_0^1 ds \int_{\mathbb{R}^6} |\nabla \bar{f} \circ \Phi_{-s}^H - \nabla \bar{f}| dx dv \right. \\ &\quad \left. + \int_0^1 ds \int_{\mathbb{R}^6} |\nabla \bar{f} \circ \Phi_{-s}^H| |\nabla \Phi_{-s}^H - \text{Id}| dx dv \right) \\ &=: \|\nabla H\|_{L^\infty} (A + B). \end{aligned}$$

We now estimate the terms A and B . By the Fundamental Theorem of Calculus,

$$\begin{aligned} A &= \int_0^1 ds \int_{\mathbb{R}^6} \left| \int_0^s \frac{d}{d\tau} \nabla \bar{f} \circ \Phi_{-\tau}^H d\tau \right| dx dv \\ &= \int_0^1 ds \int_{\mathbb{R}^6} \left| \int_0^s \nabla^2 \bar{f} \circ \Phi_{-\tau}^H \cdot J\nabla H \circ \Phi_{-\tau}^H d\tau \right| dx dv \\ &\leq \|\nabla H\|_{L^\infty} \int_0^1 ds \int_{\mathbb{R}^6} \int_0^s |\nabla^2 \bar{f}| \circ \Phi_{-\tau}^H d\tau dx dv \end{aligned}$$

By Fubini, we can rewrite the last integral above as

$$\int_0^1 ds \int_0^s d\tau \int_{\mathbb{R}^6} |\nabla^2 \bar{f}| \circ \Phi_{-\tau}^H dx dv$$

and because $\Phi_{-\tau}^H$ is measure preserving we deduce that the term above is equal to

$$\int_0^1 ds \int_0^s d\tau \int_{\mathbb{R}^6} |\nabla^2 \bar{f}| dx dv = \frac{1}{2} \|\nabla^2 \bar{f}\|_{L^1}.$$

Hence, in conclusion,

$$A \leq \frac{1}{2} \|\nabla^2 \bar{f}\|_{L^1} \|\nabla H\|_{L^\infty}.$$

For B , we want to estimate the term

$$|\nabla \bar{f} \circ \Phi_{-s}^H| |\nabla \Phi_{-s}^H - \text{Id}|.$$

Differentiating the equation in (1.6), we deduce that

$$\begin{cases} \partial_s \nabla \Phi_s^H = J\nabla^2 H(\Phi_s^H) \cdot \nabla \Phi_s^H \\ \nabla \Phi_0^H(x, v) = \text{Id}. \end{cases} \quad (2.15)$$

Thus,

$$\begin{cases} \frac{d}{ds} |\nabla \Phi_s^H| \leq \|\nabla^2 H\|_{L^\infty} |\nabla \Phi_s^H| \\ |\nabla \Phi_0^H| = 1 \end{cases} \quad (2.16)$$

and by Gronwall's inequality

$$|\nabla \Phi_s^H| \leq e^{s\|\nabla^2 H\|_{L^\infty}}. \quad (2.17)$$

Therefore,

$$\begin{aligned} |\nabla \Phi_s^H - \text{Id}| &= \left| \int_0^s \partial_\tau \nabla \Phi_\tau^H d\tau \right| \stackrel{(2.16)}{\leq} \|\nabla^2 H\|_{L^\infty} \sup_{\tau \in [0, s]} |\nabla \Phi_\tau^H| \\ &\stackrel{(2.17)}{\leq} \|\nabla^2 H\|_{L^\infty} e^{s\|\nabla^2 H\|_{L^\infty}}, \end{aligned}$$

which yields

$$B \leq \|\nabla^2 H\|_{L^\infty} e^{\|\nabla^2 H\|_{L^\infty}} \int_{\mathbb{R}^6} |\nabla \bar{f}| dx dv.$$

Combining the latter estimates, we obtain

$$\left| \int_{\mathbb{R}^6} |f_1 - \bar{f}| dx dv - \int_{\mathbb{R}^6} |g| dx dv \right| \leq C \|\nabla H\|_{L^\infty} \left(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} e^{\|\nabla^2 H\|_{L^\infty}} \right),$$

where C is a constant depending only on $\|\nabla \bar{f}\|_{L^1}$ and $\|\nabla^2 \bar{f}\|_{L^1}$. $\square$

2.5 Proof of Theorem 1.7

In this section we combine the upper and lower bounds obtained in Sections 2.2 and 2.3 with interpolation estimates to obtain a contradiction to the existence of a stationary solution $\bar{f}_1^H$ with H as in the statement of Theorem 1.7.

We begin by recalling that, by the Sobolev's embedding, given $R > 0$ and $u : B_R \rightarrow \mathbb{R}$ compactly supported,

$$\|u\|_{L^\infty(B_R)} \leq C_{n,R} \|\nabla^s u\|_{L^2(B_R)} \quad \forall s > n/2. \quad (2.18)$$

In particular, since $n = 6$, if H is as in the statement of the theorem then $\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty}$ is as small as desired provided we choose ε small enough. This allows us to apply Lemma 2.1, that combined with Lemma 2.4 yields following bound on g :

$$\|g\|_{L^1} \leq C \left(\sqrt{\mathcal{H}(\bar{f}_1^H) - \mathcal{H}(\bar{f})} + \|\nabla H\|_{L^\infty}^2 + \|\nabla H\|_{L^\infty} \|\nabla^2 H\|_{L^\infty} e^{\|\nabla^2 H\|_{L^\infty}} \right).$$

Then, using Proposition 2.2,

$$\begin{aligned} \|g\|_{L^1} &\leq C \left(\|\nabla H\|_{L^\infty} \left(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} \right)^{\frac{1}{2}} \right. \\ &\quad \left. + \|\nabla H\|_{L^\infty}^2 + \|\nabla H\|_{L^\infty} \|\nabla^2 H\|_{L^\infty} e^{\|\nabla^2 H\|_{L^\infty}} \right). \end{aligned}$$

We now use the assumption $H \in A_k$ to get

$$\begin{aligned} \|\nabla H\|_{L^1} &\leq Ck \left(\|\nabla H\|_{L^\infty} \left(\|\nabla H\|_{L^\infty} + \|\nabla^2 H\|_{L^\infty} \right)^{\frac{1}{2}} \right. \\ &\quad \left. + \|\nabla H\|_{L^\infty}^2 + \|\nabla H\|_{L^\infty} \|\nabla^2 H\|_{L^\infty} e^{\|\nabla^2 H\|_{L^\infty}} \right). \quad (2.19) \end{aligned}$$

Note that if the norms in the left hand side and in the right hand side were comparable, we would have an inequality of the form

$$\|\nabla H\|_X \leq C \|\nabla H\|_X^{3/2},$$

which is impossible when H is small enough. Thus, the next step is to use interpolation estimates to compare the different norms of ∇H appearing in (2.19). More precisely, we want to use the following interpolation estimates.

Lemma 2.5. *For any smooth compactly supported function $u : \mathbb{R}^n \rightarrow \mathbb{R}$,*

$$\|\nabla^\ell u\|_{L^2} \leq \|u\|_{L^2}^{1-\ell/m} \|\nabla^m u\|_{L^2}^{\ell/m} \quad \forall 1 \leq \ell \leq m.$$

Proof. The proof is immediate by Fourier: using Hölder inequality with the conjugate exponents m/ℓ and $(m-\ell)/\ell$, we get

$$\begin{aligned} \int |\xi|^{2\ell} |\hat{u}|^2 &= \int (|\xi|^{2\ell} |\hat{u}|^{2\ell/m}) |\hat{u}|^{2(m-\ell)/m} \leq \| |\xi|^{2\ell} |\hat{u}|^{2\ell/m} \|_{L^{m/\ell}} \| |\hat{u}|^{2(m-\ell)/m} \|_{L^{(m-\ell)/\ell}} \\ &= \left(\int |\xi|^{2m} |\hat{u}|^2 \right)^{\ell/m} \left(\int |\hat{u}|^2 \right)^{1-\ell/m}. \end{aligned}$$

Since $\|\nabla^k u\|_{L^2(\mathbb{R}^n)} = \| |\xi|^k \hat{u} \|_{L^2(\mathbb{R}^n)}$ for all $k \geq 0$, the result follows. $\square$

Since (2.19) involves L^1 and L^∞ norms, to apply Lemma 2.5 we use shall use other interpolation inequalities. More precisely, we recall the classical Nash inequality:

$$\|u\|_{L^2(\mathbb{R}^n)}^{1+2/n} \leq C_n \|u\|_{L^1(\mathbb{R}^n)}^{2/n} \|\nabla u\|_{L^2(\mathbb{R}^n)}. \quad (2.20)$$

We now set $n = 6$, and we let s be a number larger than $n/2 = 3$ to be fixed later. Applying (2.20) to $\partial_i H : \mathbb{R}^n \rightarrow \mathbb{R}$, $i = 1, \dots, n$, we get

$$\|\nabla H\|_{L^2}^{1+2/n} \leq C \|\nabla H\|_{L^1}^{2/n} \|\nabla^2 H\|_{L^2}.$$

Let us recall that, by assumption, H is supported in $B_{2\rho}$. Hence, we can apply (2.18) both with $u = \partial_i H$ and $u = \partial_{ij} H$ to get

$$\|\nabla H\|_{L^\infty} \leq C \|\nabla^{s+1} H\|_{L^2}, \quad \|\nabla^2 H\|_{L^\infty} \leq C \|\nabla^{s+2} H\|_{L^2}.$$

Note also that, by Poincaré inequality in $B_{2\rho}$, $\|\nabla^{s+1} H\|_{L^2} \leq C \|\nabla^{s+2} H\|_{L^2}$. Combining all these estimates with (2.19), we get

$$\|\nabla H\|_{L^2}^{1+2/n} \leq Ck \left(\|\nabla^{s+2} H\|_{L^2}^{3/2} + \|\nabla^{s+2} H\|_{L^2}^2 + \|\nabla^{s+2} H\|_{L^2}^2 e^{\|\nabla^{s+2} H\|_{L^2}} \right)^{2/n} \|\nabla^2 H\|_{L^2}. \quad (2.21)$$

To conclude we recall that, by assumption, $\|H\|_{W^{r,2}} \leq \varepsilon$, where $r \geq 22$, and we want to obtain a contradiction when ε is sufficiently small. To this aim, we first note that, for $s \leq r-2$,

$$\|\nabla^{s+2} H\|_{L^2} \leq \|H\|_{W^{r,2}} \leq \varepsilon \ll 1.$$

This implies that the quadratic terms in (2.21) are much smaller than the term with the power $3/2$, therefore (2.21) yields

$$\|\nabla H\|_{L^2}^{1+2/n} \leq Ck \|\nabla^{s+2} H\|_{L^2}^{3/n} \|\nabla^2 H\|_{L^2}.$$

Then we apply Lemma 2.5 with $u = \partial_i H$, $\ell = s+1$, and $m = r-1$ to get

$$\|\nabla^{s+2} H\|_{L^2} \leq C \|\nabla H\|_{L^2}^{\frac{r-s-2}{r-1}} \|\nabla^r H\|_{L^2}^{\frac{s+1}{r-1}} \leq C \varepsilon^{\frac{s+1}{r-1}} \|\nabla H\|_{L^2}^{\frac{r-s-2}{r-1}},$$

therefore

$$\|\nabla H\|_{L^2}^{1+2/n} \leq Ck \left(\|\nabla H\|_{L^2}^{\frac{r-s-2}{r-1}} \varepsilon^{\frac{s+1}{r-1}} \right)^{3/n} \|\nabla^2 H\|_{L^2}. \quad (2.22)$$

Also, by Lemma 2.5 with $u = \partial_i H$, $\ell = 1$, $m = r-1$, we have

$$\|\nabla^2 H\|_{L^2} \leq C \|\nabla H\|_{L^2}^{\frac{r-2}{r-1}} \|\nabla^r H\|_{L^2}^{\frac{1}{r-1}} \leq C \varepsilon^{\frac{1}{r-1}} \|\nabla H\|_{L^2}^{\frac{r-2}{r-1}}. \quad (2.23)$$

Thus, combining (2.22) and (2.23), we obtain

$$\|\nabla H\|_{L^2}^{1+2/n} \leq Ck\varepsilon^{\frac{1+3(s+1)/n}{r-1}} \|\nabla H\|_{L^2}^{\frac{3(r-s-2)}{n(r-1)} + \frac{r-2}{r-1}}.$$

We finally choose s . Since s is any exponent larger than $n/2 = 3$ and less than $r - 2 \geq 20$, we fix $s = 4$. Then, the inequality above becomes

$$\|\nabla H\|_{L^2}^{4/3} \leq Ck\varepsilon^{\frac{1+3(s+1)/n}{r-1}} \|\nabla H\|_{L^2}^{\frac{r-6}{2(r-1)} + \frac{r-2}{r-1}} = Ck\varepsilon^{\frac{7}{2(r-1)}} \|\nabla H\|_{L^2}^{\frac{3r-10}{2(r-1)}}.$$

Since $r \geq 22$ by assumption, we see that $\frac{3r-10}{2(r-1)} \geq 4/3$, thus we obtain

$$1 \leq Ck\varepsilon^{\frac{7}{2(r-1)}} \|\nabla H\|_{L^2}^{\frac{3r-10}{2(r-1)} - \frac{4}{3}} \leq Ck\varepsilon^{\frac{7}{2(r-1)} + \frac{3r-10}{2(r-1)} - \frac{4}{3}} = Ck\varepsilon^{1/6},$$

which is false for ε small enough. This shows the desired contradiction and completes the proof. $\square$

Acknowledgments: The author is grateful to Clément Mouhot for proposing to me this beautiful problem and for interesting discussions. Also, I wish to thank Pierre Raphaël for useful comments during the preparation of this manuscript. The author would also like to acknowledge the L'Oréal Foundation for partially supporting this project by awarding the L'Oréal-UNESCO *For Women in Science France fellowship*.

References

- [1] L. Ambrosio, M. Colombo, and A. Figalli. On the Lagrangian structure of transport equations: the Vlasov-Poisson system. Preprint, 2015.
- [2] V. I. Arnold. Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits. *Ann. Inst. Fourier (Grenoble)*, 16:319–361, 1966.
- [3] A. A. Arseniev. Existence in the large of a weak solution of Vlasov's system of equations. *Ž. Vyčisl. Mat. i Mat. Fiz.*, 15:136–147, 276, 1975.
- [4] J. Binney and S. Tremaine. *Galactic Dynamics*. Princeton: Princeton University Press, 1987.
- [5] A. Bohun, F. Bouchut, and G. Crippa. Lagrangian solutions to the Vlasov-Poisson equation with L^1 density. *J. Differential Equations* 260(4):3576–3597, 2016.
- [6] A. Choffrut, V. Šverák, *Local structure of the set of steady-state solutions to the 2D incompressible Euler equations*. *Geom. Funct. Anal.* 22(1):136–201, 2012.
- [7] R. J. DiPerna and P.-L. Lions. Solutions globales d'équations du type Vlasov-Poisson. (French) [Global solutions of Vlasov-Poisson type equations] *C. R. Acad. Sci. Paris Sér. I Math.*, 307:655–658, 1988.
- [8] R. J. DiPerna and P.-L. Lions. Global weak solutions of kinetic equations. *Rend. Sem. Mat. Univ. Politec. Torino*, 46:259–288, 1988.
- [9] R. J. DiPerna & P.-L. Lions. Global weak solutions of Vlasov-Maxwell systems. *Comm. Pure Appl. Math.*, 42:729–757, 1989.

- [10] A. M. Fridman, V. L. Polyachenko. *Physics of Gravitating Systems I*. New York: Springer-Verlag 1984
- [11] E. Horst and R. Hunze. Weak solutions of the initial value problem for the unmodified nonlinear Vlasov equation. *Math. Methods Appl. Sci.* 6:262–279, 1984.
- [12] S. V. Iordanskii. The Cauchy problem for the kinetic equation of plasma. *Trudy Mat. Inst. Steklov.*, 60:181–194, 1961.
- [13] J. Jeans. On the theory of star-streaming and the structure of the universe. *Mon. Not. R. Astron. Soc.* 76:70–84, 1915, and 76:552–567, 1916.
- [14] M. Lemou. Extended rearrangement inequalities and applications to some quantitative stability results. Preprint, 2015.
- [15] M. Lemou, F. Méhats, and P. Raphaël. Orbital stability of spherical galactic models. *Invent. Math.*, 187(1):145–194, 2012.
- [16] P.-L. Lions and B. Perthame. Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system. *Invent. Math.*, 105:415–430, 1991.
- [17] C. Mouhot. Stabilité orbitale pour le système de Vlasov-Poisson gravitationnel (d’après Lemou-Méhats-Raphaël, Guo, Lin, Rein et al.). (French) [Orbital stability for the gravitational Vlasov-Poisson system (after Lemou-Méhats-Raphaël, Guo, Lin, Rein et al.)] Séminaire Bourbaki. Vol. 2011/2012. Exposés 1043-1058. *Astérisque* No. 352 (2013), Exp. No. 1044, vii, 35-82.
- [18] K. Pfaffelmoser. Global classical solutions of the Vlasov-Poisson system in three dimensions for general initial data. *J. Differential Equations*, 95(2):281–303, 1992.
- [19] J. Schaeffer. Global existence of smooth solutions to the Vlasov-Poisson system in three dimensions. *Comm. Partial Differential Equations*, 16(8-9):1313–1335, 1991.
- [20] S. Ukai and T. Okabe. On classical solutions in the large in time of two-dimensional Vlasov’s equation. *Osaka J. Math.*, 15:245–261, 1978.
- [21] A. A. Vlasov. *Zh. Eksper. Teor. Fiz.* 8:291, 1938.
- [22] A. A. Vlasov. Vlasov equation and plasma dispersion relation. *J. Phys. (U.S.S.R.)* 9:25, 1945.